
Figure 1


## TOPICAL INDUCERS OF LANGERHANS CELL MIGRATION



Cell Number/Lymph Node x 10<sup>-4</sup>

Figure 2

## DENDRITIC CELLS In LYMPH NODE after SKIN PAINTING



## LYMPH NODE CELLULARITY after SKIN PAINTING




Figure 4





Figure 5

and DBP Induces Protective Tumor Immunity Intravaginal Application of Peptide








FIGURE CELEBER

Topical Application of DibutyIphthalate Analogs Lymph Node Dendritic Cells Induced by




Quantitative Detection of Peptide: MHC Complexes on Langerhans Cells in vitro




Complexes on Langerhans Cells in vitro

Ultrasound Application of Soluble Peptide followed by Migration Inducer Antigen-bearing Dendritic Cells in Lymph Nodes after




Injection of Soluble Protein or Peptide followed by Migration Inducer Antigen-bearing Dendritic Cells in Lymph Nodes after



**MHC Class II** 


Injection of Soluble Peptide with or without subsequent Migration Inducer Antigen-bearing Dendritic Cells in Lymph Nodes after




...


Fig. 12

Tumor-specific Immunity induced by a Single Injection of a Tumor Protein followed by a Topical Migration Inducer





## Adjuvant Effects f Inducing Dendritic Cell Migration/Maturation



Antigen+ Dendritic Cells

504  $\longrightarrow$  22,239

2,520  $\longrightarrow$  33,673

Lymphocytes  $6 \times 10^6 \longrightarrow 8.6 \times 10^6$